

Gene dosage defects in primary immunodeficiencies and related disorders: a pilot study

Lisa Dyer^{1,2}, Xia Li^{1,2}, James Denton¹, Brittany Jones¹, Emily Liston¹, Danielle Hilton¹, Abhinav Mathur¹, Kejian Zhang^{1,2}, C. Alexander Valencia^{1,2}

¹Division of Human Genetics, Cincinnati Children's Hospital Medical Center, Cincinnati, OH 45229, USA.

²Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH 45267, USA.

Correspondence to: Dr. C. Alexander Valencia, Division of Human Genetics, Cincinnati Children's Hospital Medical Center, Cincinnati, OH 45229, USA. E-mail: avalenci2001@gmail.com

How to cite this article: Dyer L, Li X, Denton J, Jones B, Liston E, Hilton D, Mathur A, Zhang K, Valencia CA. Gene dosage defects in primary immunodeficiencies and related disorders: a pilot study. *J Transl Genet Genom* 2017;1:23-7.

Article history: Received: 11 Dec 2017 First Decision: 18 Dec 2017 Revised: 20 Dec 2017 Accepted: 20 Dec 2017 Published: 25 Dec 2017

TO THE EDITOR

Detection of structural variations or copy number variants (CNVs) is a critical component of diagnosing many genetic disorders especially when the causative mutation(s) cannot be identified in dominant, recessive and X-linked disorders by Sanger or next generation sequencing. Fluorescence *in situ* hybridization, whole genome copy number microarrays, next generation sequencing, Sanger sequencing, multiplex ligation-dependent probe amplification, and quantitative real-time polymerase chain reaction (PCR) may miss potential pathogenic structural variations that range in size larger than 500 bp or includes at least one exon. Moreover, CNV calls extracted from whole exome sequencing remain a challenge for clinical genetic testing laboratories due to high false positive rates. To overcome these technical challenges, we designed a custom high-resolution targeted exon-centric microarray (ECM) that has 1508 disease-causing genes including complementary genes to our

laboratory's immunodeficiency focused next-generation sequencing (NGS) panels. The combination of targeting rare immunodeficiency genes with large case load of clinical NGS testing for rare immunological and related disorders allowed for the opportunity to discover novel CNVs and expand our understanding of disease mechanisms in this group of genetically heterogeneous disorders. We have been routinely performing the ECM as a clinical assay and herein report a number of novel CNVs detected in this group of disorders for which intragenic deletions are not a commonly reported etiology. We followed the diagnostic algorithm illustrated in [Figure 1](#), to increase the diagnostic yield in patients in this group of disorders, but albeit is not limited to these disorders and is recommended for Mendelian disorders because the mutation spectrum of many genes remain understudied and deletions and duplications may represent a significant percentage of pathogenic changes.

Clinical array comparative genomic hybridization was



This is an open access article licensed under the terms of Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original author is credited and the new creations are licensed under the identical terms.

For reprints contact: service@oaepublish.com

Quick Response Code:






Figure 1: CNV testing algorithm for Mendelian disorders. The diagram demonstrates the diagnostic work-up of a Mendelian disorder using next generation sequencing as the first step followed by deletion/duplication testing to find an allele or second allele. CNV: copy number variant; NGS: next-generation sequencing

performed on 1048 patient samples referred to the Molecular Genetics Laboratory at Cincinnati Children's Hospital Medical Center (CCHMC). ECM analysis was performed on only the gene(s) of interest. Reference DNA was obtained from Promega (Madison, WI). This study was approved by the Institutional Review Board at CCHMC.

The custom 4X180K microarray was developed on the Agilent Technologies (Santa Clara, CA) platform using the GeneEfficiency service (Oxford Gene Technology, Oxford, UK). OGT used proprietary ink-jet *in situ* printer technology (IJISS) developed by Rosetta InPharmatics (Kirkland, WA) and Agilent Technologies that allows *in situ* synthesis of long oligonucleotides. The probes were 60 bp in length and annotated against NCBI build 37 (UCSC hg19, March 2006). Included were 364,591 probes that targeted 1508 genes (exons and introns), up to 300 bp flanking the exon/intron boundaries and backbone coverage.

DNA was extracted from whole blood collected in EDTA (purple-top) collection tubes using the Puregene DNA extraction kit (Qiagen, Valencia, CA) or the Chemagic magnetic separation module I (PerkinElmer, Baesweiler, Germany) according to the manufacturer's recommendations. Comparative genomic hybridization was performed following the manufacturer's protocol (Agilent Technologies, Santa Clara, CA). The reference DNA was used from two pools (male and

female) from normal individuals, run as a same-sex control. DNA was sonicated using a Branson Sonifier 450 with cup horn (Danbury, CT) and visualized on a two-percent agarose gel prior to labeling, as a quality control measure. Each patient and reference DNA was labeled with Cy3 and Cy5 9-mer primers, respectively. Purification of labeled products, hybridization, and post-wash of the array was carried out according to Agilent's recommendation and with their proprietary solutions. Microarray slides were scanned with Innosys Innoscan 710 and raw data was generated with extraction software (Chicago, Illinois).

CytoSure Interpret software version 4.4 (OGT) was used for analysis of microarray data (referred to as CytoSure). The program uses the circular binary segmentation algorithm to generate segments along the chromosomes that have similar copy number relative to reference chromosome. Averaging of the segments is with median value of all segments on a chromosome as the baseline. Deletion or duplication CNV calls were made using the log₂ ratio of each segment with a minimum of 4 unique and consequent probes. The threshold factor for deletions was set as a log₂ ratio of -0.6 that is less stringent than the theoretical log₂ score of -1 for heterozygous deletions [heterozygous deletion log₂(1/2) = -1; no change in allele number log₂(2/2) = 0; heterozygous duplication log₂(3/2) = 0.59; hemizygous deletion log₂(0/1) = -1; homozygous deletion log₂(0/2) = -2]. The software uses the standard deviation of the log₂ ratio to calculate a deviation log ratio (DLR), which is used as a quality control check. A DLR of 0.08-0.19 is acceptable, 0.20-0.29 is borderline, and ≥ 0.30 is rejected. The DLR for all microarrays shown was scored by this scale. Data was analyzed only for the gene(s) ordered for testing.

We used the high-resolution ECM to process 1048 of samples that were reflex testing from negative or equivocal targeted NGS panel testing or ordered for single gene assessment. We identified 23 CNVs in rare immunodeficiency and related genes [Table 1]. Of the 23 CNVs detected, 21 deletions were identified. Significantly, of these 23 CNVs, 11 (48%) of them were novel, for which intragenic deletions and duplications are not a commonly reported etiology, either due to the low mutation burden of these events or the lack of recognition by an appropriate testing methodology. Six of the 23 CNVs in immunodeficiency and related genes were identified in dominantly inherited genes (26%), 7 hemizygous CNVs were identified in male patients in X-linked recessive genes (30%), 5 CNVs were identified as the second allele in 3 autosomal recessive genes (17%), and surprisingly,

Table 1: List of patients with CNVs identified by exon-centric microarray testing

Patient	Gene	OMIM phenotype: inheritance	CNV coordinates (min)	Min size (Max size)	Min region affected	Protein domain affected	Sequencing results	Reference
1	<i>FAS</i>	Autoimmune lymphoproliferative syndrome: AD	arr[hg19] 10q23.31 (90,770,398-90,777,298) x1	6.9 Kb (53.59 Kb)	exons 6-9	Death-like	No variant identified	This study
2	<i>FAS</i>	Autoimmune lymphoproliferative syndrome: AD	arr[hg19] 10q23.31 (90,748,451-91,780,017) x1	1.03 Mb (1.19 Mb)	whole gene		No variant identified	This study
3 [^]	<i>FAS</i>	Autoimmune lymphoproliferative syndrome: AD	arr[hg19] 10q23.21(90774157-90774381)x0	224bp (403bp)	exon 9	Death-like	No variant identified	This study
4	<i>RPS26</i>	Diamond-Blackfan anemia: AD	arr[hg19] 12q13.2q13.3 (56,175,892-56,691,772) x1	518.88 Kb (688.55 Kb)	whole gene		N/A	[6]
5	<i>RPL35A</i>	Diamond-Blackfan anemia: AD	arr[hg19] 3q29 (197,456,033-197,844,960) x1	388.93 Kb (855.23 Kb)	whole gene		N/A	[6]
6	<i>TERT</i>	Dyskeratosis congenita: AD	arr[hg19] 5p15.33 (1,226,629-1,286,091)x1	59.46 Kb (60.16 Kb)	exon 3-16	Reverse transcriptase	No variant identified	This study
7 [*]	<i>WAS</i>	Wiskot-Aldrich syndrome: XLR	arr[hg19] Xp11.23 (48,533,049-48,542,539) x0	9.49 Kb (147.2 Kb)	exon 1	EVH1	No variant identified	This study
8 [*]	<i>WAS</i>	Wiskot-Aldrich syndrome: XLR	arr[hg19] Xp11.23 (48,533,302-48,613,923) x0	80.62 Kb (110.45 Kb)	whole gene		No variant identified	[7]
9	<i>WAS</i>	Wiskot-Aldrich syndrome: XLR	arr[hg19] Xp11.23 (48,533,049-48,550,909) x0	17.86 Kb (220.16 Kb)	whole gene		No variant identified	[7]
10	<i>WAS</i>	Wiskot-Aldrich syndrome: XLR	arr[hg19] Xp11.23 (48,533,049-48,551,398) x1	18.35 Kb (282.83 Kb)	whole gene		No variant identified	[7]
11	<i>SH2D1A</i>	X-linked recessive lymphoproliferative disorder: XLR	arr[hg19] Xq25 (123,481,887-123,501,318) x0	19.43 Kb (20.58 Kb)	exon 2	SH2	No variant identified	[8,9]
12	<i>XIAP</i>	X-linked recessive lymphoproliferative disorder: XLR	arr[hg19] Xq25 (123,007,906-123,021,550) x0	13.64 Kb (15.61 Kb)	exon 2	Baculoviral inhibition of apoptosis protein repeat	No variant identified	This study
13	<i>MAGT1</i>	Immunodeficiency: XLR	arr[hg19]Xq21.1 (77,088,396-77,150,089) x0	63 Kb (63 Kb)	Exons 2-7	Oligosaccharyl transferase complex	No variant identified	This study
14	<i>IL7R</i>	Severe combined immunodeficiency, T-cell -, B-cell +: AR	arr[hg19] 5p13.2 (35,867,205-35,867,604) x1	399 bp (3.94 Kb)	exon 3		c.669_688del20 (p.S223fs)	[10]
15	<i>FANCA</i>	Fanconi Anemia: AR	arr[hg19] 16q24.3 (89,880,335-89,928,824) x0	48.49 Kb (107.96 Kb)	exons 1-3		No variant identified	[11]
16	<i>FANCA</i>	Fanconi Anemia: AR	arr[hg19] 16p24.3 (89,874,512-89,884,589) x1	10.08kb (56.89 Kb)	exons 1-6		c.862G>T(p. E288X)	[12]
17	<i>FANCA</i>	Fanconi Anemia: AR	arr[hg19] 16q24.3 (89,880,335-89,928,824) x1	48.49 Kb (107.96 Kb)	exons 1-3		c.3918dupT(p. Q1307fs)	[11]
18	<i>FANCC</i>	Fanconi Anemia: AR	arr[hg19] 9q22.32 (98,007,056-98,011,765) x1	4.71 Kb (5.77 Kb)	exons 2-3		c.1642C>T(p. R548*)	[13,14]
19	<i>RAD51C</i>	Fanconi Anemia: AR	arr[hg19] 17q22 (56,765,347-56,790,584) x3	25.24 Kb (156.62 Kb)	exons 1-5	DNA recombination and repair, ATP binding domain	No variant identified	This study

20	<i>LRBA</i>	Immunodeficiency: AR	arr[hg19] 4q31.3 (151,198,604-151,199,333) x0	729 bp (7.59 Kb)	exon 57	WD40 repeat	No variant identified	This study
21 [^]	<i>STXBP2</i>	Hemophagocytic lymphohistiocytosis: AR	arr[hg19] 19p13.2 (7,706,766-7,781,506) x0	74.74 Kb (426.44 Kb)	exons 8-19	Sec-like protein	No variant identified	This study
22	<i>RAB27A</i>	Griscelli syndrome, type 2: AR	arr[hg19] 15q21.3 (55,499,521-55,562,873) x1	63.35 Kb (67.75 Kb)	exons 1-5	GTPase	No variant identified	[15]
23	<i>RAB27A</i>	Griscelli syndrome, type 2: AR	arr[hg19] 15q21.3 (55,514,892-55,550,411) x3	35.52 Kb (40.86 Kb)	exons 2-5	GTPase	No variant identified	This study

*Maternal inheritance confirmed; [^]parental carrier status confirmed. CNV: copy number variant; AD: autosomal dominant; AR: autosomal recessive; XLR: X-linked recessive

3 homozygous deletions were identified (13%).

Meaningfully, we identified 3 novel CNVs in the *FAS* gene. Heterozygous mutations in the *FAS* gene cause autoimmune lymphoproliferative syndrome (ALPS), an autosomal dominant disorder, where only three CNVs have been previously described^[1-4]. In patient 1, we identified a 6.9 Kb heterozygous deletion that encompassed exons 6-9 and included the intracellular domain and the death domain of the *FAS* protein. Single gene CNV assessment of *FAS* was ordered for this patient after negative sequencing of *FAS*, *FASLG*, and *CASP10* genes. This patient's phenotype included splenomegaly, thrombocytopenia, history of acute cerebellitis and failure to thrive. Flow cytometry studies from an external laboratory identified an increased population of CD4/CD8 double negative T cells and reduced *FAS* protein expression. Similarly, in patient 2 and 3, we identified a 1.03 Mb whole gene heterozygous deletion and a 224 bp homozygous deletion in exon 9, respectively. ECM testing was ordered as reflex testing from Sanger sequencing in all 3 patients. The identification of the *FAS* gene deletions in these patients confirmed a diagnosis of ALPS. In addition, the CNV mutation spectrum of *FAS* was doubled, showing that ECM is a powerful tool for CNV discovery in rare Mendelian disorders.

Other rare hematology and immunodeficiency genes for which novel CNVs were identified included *TERT*, *WAS*, *XIAP*, *MAGT1*, *RAD51C*, *LRBA*, *STXBP2*, and *RAB27A*. Deletions in X-linked genes showed high clinical utility in males due to their hemizygous status. In several cases, a lack of amplification by PCR suggested a deletion, but results were inconclusive in clinical interpretation due to lack of direct evidence and insufficient controls. *WAS*-related disorders are X-linked recessive and include Wiskott-Aldrich syndrome, X-linked thrombocytopenia, and X-linked congenital neutropenia. Sequencing of the *WAS*

gene identifies up to 95% of pathogenic variants and the remaining 5% of variants are detectable by deletion/duplication testing and 15 gross deletions have been reported. We identified 3 CNVs in 3 male patients involving the *WAS* gene. In patient 7, we identified a previously unreported 9.49 Kb intragenic deletion of exon 1, and in patients 8, 9 and 10, we identified previously reported whole gene deletions in *WAS* of size 80.62 Kb, 17.86 Kb, and 18.35 Kb, respectively. Autosomal dominant mutations in *TERT* cause dyskeratosis congenital, a rare multisystem disorder caused by defective telomere maintenance. We identified a novel 59.46 Kb heterozygous deletion of exons 3-16 and included the reverse transcriptase domain of the *TERT* protein, the catalytic component of the enzyme. To date, only one CNV in *TERT* had been identified in a family with adult-onset pulmonary fibrosis^[4]. In addition, the *LRBA* protein is involved in regulating recycling of the cytotoxic T lymphocytes-associated protein 4, which helps regulate autoimmune responses. Biallelic deleterious mutations in *LRBA* have been associated with common variable immunodeficiency type 8 and to date, only 6 mutations have been identified in *LRBA*. The biological role of *LRBA* is currently being elucidated and recent publications have suggested targeted therapies for patients with *LRBA* deficiency^[5]. We identified a novel 729 bp homozygous deletion of exon 57 of *LRBA* in a patient from consanguineous parents.

In our cohort, we recognized a high diagnostic yield for the ECM assays for immunodeficiencies and related conditions. However, functional studies will be necessary to confirm the causative relationship to disease. This report emphasizes the importance of appropriate ECM testing for immunodeficiency disorders and helps further define gene mutation spectra and our understanding of disease mechanisms in this understudied group of genetically heterogeneous disorders.

DECLARATIONS

Authors' contributions

Conceptualized and initiated the study, collected, analyzed, and interpreted the data, wrote and reviewed the manuscript: C.A. Valencia

Analyzed and interpreted the data, wrote and reviewed the manuscript: L. Dyer, X. Li, K. Zhang

Developed the test and reviewed the manuscript: A. Mathur, J. Denton

Performed the experiments, collected, analyzed and interpreted the data and reviewed the manuscript: B. Jones, E. Liston, D. Hilton

Approved the final manuscript as submitted and agree to be accountable for all aspects of the work: all authors

Financial support and sponsorship

The study was supported by the clinical molecular genetics laboratory at CCHMC.

Conflicts of interest

The authors declare that they have no relevant conflicts of interest.

Patient consent

Not applicable.

Ethics approval

This study was approved by the Institutional Review Board at Cincinnati Children's Hospital Medical Center.

REFERENCES

- van der Werff ten Bosch JE, Demanet C, Balduck N, Bakkus MH, De Raeye H, Desprechins B, Otten J, Thielemans K. The use of the anti-malaria drug Fansidar (pyrimethamine and sulphadoxine) in the treatment of a patient with autoimmune lymphoproliferative syndrome and Fas deficiency. *Br J Haematol* 1998;102:578-81.
- Magerus-Chatinet A, Neven B, Stolzenberg MC, Daussey C, Arkwright PD, Lanzarotti N, Schaffner C, Cluet-Dennetiere S, Haerynck F, Michel G, Bole-Feysot C, Zarhrate M, Radford-Weiss I, Romana SP, Picard C, Fischer A, Rieux-Laucat F. Onset of autoimmune lymphoproliferative syndrome (ALPS) in humans as a consequence of genetic defect accumulation. *J Clin Invest* 2011;121:106-12.
- Pensati L, Costanzo A, Ianni A, Accapezzato D, Iorio R, Natoli G, Nisini R, Almerighi C, Balsano C, Vajro P, Vegnente A, Levrero M. Fas/Apo1 mutations and autoimmune lymphoproliferative syndrome in a patient with type 2 autoimmune hepatitis. *Gastroenterology* 1997;113:1384-9.
- Tsakiri KD, Cronkhite JT, Kuan PJ, Xing C, Raghu G, Weissler JC, Rosenblatt RL, Shay JW, Garcia CK. Adult-onset pulmonary fibrosis caused by mutations in telomerase. *Proc Natl Acad Sci U S A* 2007;104:7552-7.
- Lo B, Zhang K, Lu W, Zheng L, Zhang Q, Kanellopoulou C, Zhang Y, Liu Z, Fritz JM, Marsh R, Husami A, Kissell D, Nortman S, Chaturvedi V, Haines H, Young LR, Mo J, Filipovich AH, Bleesing JJ, Mustillo P, Stephens M, Rueda CM, Choungnet CA, Hoebe K, McElwee J, Hughes JD, Karakoc-Aydiner E, Matthews HF, Price S, Su HC, Rao VK, Lenardo MJ, Jordan MB. Patients with LRBA deficiency show CTLA4 loss and immune dysregulation responsive to abatacept therapy. *Science* 2015;349:436-40.
- Quarello P, Garelli E, Brusco A, Carando A, Mancini C, Pappi P, Vinti L, Svahn J, Dianzani I, Ramenghi U. High frequency of ribosomal protein gene deletions in Italian Diamond-Blackfan anemia patients detected by multiplex ligation-dependent probe amplification assay. *Haematologica* 2012;97:1813-7.
- Imai K, Morio T, Zhu Y, Jin Y, Itoh S, Kajiwarra M, Yata J-I, Mizutani S, Ochs HD, Nonoyama S. Clinical course of patients with WASP gene mutations. *Blood* 2004;103:456-64.
- Yin L, Ferrand V, Lavoué MF, Hayoz D, Philippe N, Souillet G, Seri M, Giacchino R, Castagnola E, Hodgson S, Sylla BS, Romeo G. SH2D1A mutation analysis for diagnosis of XLP in typical and atypical patients. *Hum Genet* 1999;105:501-5.
- Sumegi J, Huang D, Lanyi A, Davis JD, Seemayer TA, Maeda A, Klein G, Seri M, Wakiguchi H, Purtilo DT, Gross TG. Correlation of mutations of the SH2D1A gene and Epstein-Barr virus infection with clinical phenotype and outcome in X-linked lymphoproliferative disease. *Blood* 2000;96:3118-25.
- Bayer DK, Martinez CA, Sorte HS, Forbes LR, Demmler-Harrison GJ, Hanson IC, Pearson NM, Noroski LM, Zaki SR, Bellini WJ, Leduc MS, Yang Y, Eng CM, Patel A, Rodningen OK, Muzny DM, Gibbs RA, Campbell IM, Shaw CA, Baker MW, Zhang V, Lupski JR, Orange JS, Seeborg FO, Stray-Pedersen A. Vaccine-associated varicella and rubella infections in severe combined immunodeficiency with isolated CD4 lymphocytopenia and mutations in IL7R detected by tandem whole exome sequencing and chromosomal microarray. *Clin Exp Immunol* 2014;178:459-69.
- Castella M, Pujol R, Callén E, Trujillo JP, Casado JA, Gille H, Lach FP, Auerbach AD, Schindler D, Benítez J, Porto B, Ferro T, Muñoz A, Sevilla J, Madero L, Cela E, Beléndez C, de Heredia CD, Olivé T, de Toledo JS, Badell I, Torrent M, Estella J, Dasí A, Rodríguez-Villa A, Gómez P, Barbot J, Tapia M, Molinés A, Figuera A, Bueren JA, Surrallés J. Origin, functional role, and clinical impact of Fanconi anemia FANCA mutations. *Blood* 2011;117:3759-69.
- Levrano O, Diotti R, Pujara K, Batish SD, Hanenberg H, Auerbach AD. Spectrum of sequence variations in the FANCA gene: an International Fanconi Anemia Registry (IFAR) study. *Hum Mutat* 2005;25:142-9.
- Lo ten Foe JR, Barel MT, Thuss P, Digweed M, Arwert F, Joenje H. Sequence variations in the Fanconi anaemia gene, FAC: pathogenicity of 1806insA and R548X and recognition of D195V as a polymorphic variant. *Hum Genet* 1996;98:522-3.
- Hartmann L, Neveling K, Borkens S, Schneider H, Freund M, Grassman E, Theiss S, Wawer A, Burdach S, Auerbach AD, Schindler D, Hanenberg H, Schaal H. Correct mRNA processing at a mutant TT splice donor in FANCC ameliorates the clinical phenotype in patients and is enhanced by delivery of suppressor U1 snRNAs. *Am J Hum Genet* 2010;87:480-93.
- Anikster Y, Huizing M, Anderson PD, Fitzpatrick DL, Klar A, Gross-Kieselstein E, Berkun Y, Shazberg G, Gahl WA, Hurvitz H. Evidence that Griscelli syndrome with neurological involvement is caused by mutations in RAB27A, not MYO5A. *Am J Hum Genet* 2002;71:407-14.